

# Stochastic Processes



**Week 04 (Version 1.2)**

**Poisson Processes**

**Point Process**

Hamid R. Rabiee

Fall 2024

# Outline of Week 04 Lectures

- Poisson Process
- Point Process

# Recall: Binomial Distribution and its relation to Poisson Distribution

Binomial Distribution:  $X \sim B(n, p)$

probability of exactly  $k$  success in  $n$  trials:

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

$$B(n, p) \xrightarrow[\substack{n \rightarrow \infty \\ np \text{ remains constant}}]{} \text{Poisson}(np)$$



# Poisson Processes

- Recall: Binomial and Poisson distributions:  
Both distributions can be used to model the number of occurrences of some event.
- Recall: **Poisson arrivals** are the limiting behavior of **Binomial random variables**. (Refer to Poisson approximation of Binomial random variables in your textbook):

$$P \left\{ \begin{array}{l} \text{"} k \text{ arrivals occur in an} \\ \text{interval of duration } \Delta \text{"} \end{array} \right\} = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

Where:

$$\lambda = np = \mu T \cdot \frac{\Delta}{T} = \mu \Delta$$



# Poisson Processes

It follows that:

$$P \left\{ \begin{array}{l} "k \text{ arrivals occur in an} \\ \text{interval of duration } 2\Delta" \end{array} \right\} = e^{-2\lambda} \frac{(2\lambda)^k}{k!}, \quad k = 0, 1, 2, \dots,$$

since in that case:

$$np_1 = \mu T \cdot \frac{2\Delta}{T} = 2\mu\Delta = 2\lambda.$$

# Poisson Processes

- **Poisson arrivals** over an interval form a Poisson random variable whose parameter depends on the duration of that interval.
- Moreover because of the Bernoulli nature of the underlying basic random arrivals, **events over nonoverlapping intervals are independent.**
- We shall use these **two key observations** to define a Poisson process formally.

# Poisson Process

**Definition:**  $X(t) = n(0, t)$  represents a Poisson process if:

(i) the number of arrivals  $n(t_1, t_2)$  in an interval  $(t_1, t_2)$  of length  $t = t_2 - t_1$  is a Poisson random variable with parameter  $\lambda t$ .

Thus:

$$P\{n(t_1, t_2) = k\} = e^{-\lambda t} \frac{(\lambda t)^k}{k!} \quad k = 0, 1, 2, \dots, t = t_2 - t_1$$

And:

# Poisson Processes

(ii) If the intervals  $(t_1, t_2)$  and  $(t_3, t_4)$  are nonoverlapping, then the random variables  $n(t_1, t_2)$  and  $n(t_3, t_4)$  are independent.

Since  $n(0, t) \sim P(\lambda t)$  we have:

$$E[X(t)] = E[n(0, t)] = \lambda t$$

And:

$$E[X^2(t)] = E[n^2(0, t)] = \lambda t + \lambda^2 t^2$$



# Poisson Processes

Autocorrelation function  $R_{xx}(t_1, t_2)$ :

$$R_{xx}(t_1, t_2) = E[X(t_1)X(t_2)]$$

$$X(t_1) = n(0, t_1) \text{ and } X(t_2) = n(0, t_2)$$

To determine the autocorrelation function  $R_{xx}(t_1, t_2)$  let  $t_2 > t_1$  then from (ii) above  $n(0, t_1)$  and  $n(t_1, t_2)$  are **independent Poisson random variables** with parameters  $\lambda t_1$  and  $\lambda(t_2 - t_1)$  respectively.

Thus:

$$E[n(0, t_1)n(t_1, t_2)] = E[n(0, t_1)]E[n(t_1, t_2)] = \lambda^2 t_1 (t_2 - t_1)$$

But:

$$n(t_1, t_2) = n(0, t_2) - n(0, t_1) = X(t_2) - X(t_1)$$

And:

$$E[X(t_1)\{X(t_2) - X(t_1)\}] = R_{xx}(t_1, t_2) - E[X^2(t_1)]$$

We obtain:

$$R_{xx}(t_1, t_2) = \lambda^2 t_1 (t_2 - t_1) + E[X^2(t_1)] = \lambda t_1 + \lambda^2 t_1 t_2$$
$$t_2 \geq t_1$$

Similarly, for  $t_1 > t_2$ :

$$R_{xx}(t_1, t_2) = \lambda t_2 + \lambda^2 t_1 t_2$$

Thus:

$$R_{xx}(t_1, t_2) = \lambda^2 t_1 t_2 + \lambda \min(t_1, t_2)$$

## Poisson Distribution vs Poisson Processes

**Poisson Distribution:** A discrete probability distribution that expresses the probability of a given number of events occurring in a fixed interval of time or space.

**Characteristics:** It assumes that these events occur with a known constant mean rate and independently of the time since the last event.

**Example:** The number of emails received in an hour can be modeled using a Poisson distribution if emails arrive independently and at a constant average rate.


## Poisson Distribution vs Poisson Processes

**Poisson Process:** A stochastic process that models a series of events occurring randomly over time or space.

**Characteristics:** It describes the occurrence of events that happen independently and at a constant average rate. The time between consecutive events follows an exponential distribution.

**Example:** The arrival of customers at a bank can be modeled as a Poisson process if the arrivals are independent and occur at a constant average rate.

## Example:



(Derivative as a LTI system)

Then:

$$\mu_{x'}(t) = \frac{d\mu_x(t)}{dt} = \frac{d\lambda t}{dt} = \lambda, \quad \text{a constant}$$

And:

$$\begin{aligned} R_{xx'}(t_1, t_2) &= \frac{\partial R_{xx}(t_1, t_2)}{\partial t_2} = \begin{cases} \lambda^2 t_1 & t_1 \leq t_2 \\ \lambda^2 t_1 + \lambda & t_1 > t_2 \end{cases} \\ &= \lambda^2 t_1 + \lambda U(t_1 - t_2) \end{aligned}$$

And:

$$R_{xx'}(t_1, t_2) = \frac{\partial R_{xx'}(t_1, t_2)}{\partial t_1} = \lambda^2 + \lambda \delta(t_1 - t_2).$$

# Poisson Processes

Notice that:

- The Poisson process  $X(t)$  *does not* represent a wide sense stationary process.
- Although  $X(t)$  *does not* represent a **wide sense stationary process**, its derivative  $X'(t)$  *does* represent a **wide sense stationary process**.

# Poisson Processes

Since  $X'(t)$  is a **wide sense stationary process**; nonstationary inputs to a LTI systems *can* lead to **wide sense stationary** outputs, an interesting observation!


- **Sum of Poisson Processes:**

If  $X_1(t)$  and  $X_2(t)$  represent two **independent Poisson processes**, then their sum  $X_1(t) + X_2(t)$  is also a **Poisson process** with parameter  $(\lambda_1 + \lambda_2)t$ . (Follows from the definition of the Poisson process in (i) and (ii)).

# Poisson Processes

## Random selection of Poisson Points:

Let  $t_1, t_2, \dots, t_i, \dots$  represent random arrival points associated with a Poisson process  $X(t)$  with parameter  $\lambda t$ , and associated with each arrival point, define an independent Bernoulli random variable  $N_i$ , where:



$$P(N_i = 1) = p, \quad P(N_i = 0) = q = 1 - p.$$



# Poisson Processes

Define the processes:

$$Y(t) = \sum_{i=1}^{X(t)} N_i \quad ; \quad Z(t) = \sum_{i=1}^{X(t)} (1 - N_i) = X(t) - Y(t)$$

We claim that both  $Y(t)$  and  $Z(t)$  are **independent Poisson processes** with parameters  $\lambda pt$  and  $\lambda qt$ , respectively, where  $q = 1 - p$ .  
When  $X(t)$  is a Poisson process with parameter  $\lambda t$ .

# Poisson Processes

**Proof:**

$$Y(t) = \sum_{n=k}^{\infty} P\{Y(t) = k \mid X(t) = n\} P\{X(t) = n\}.$$

But given  $X(t) = n$ , we have  $Y(t) = \sum_{i=1}^n N_i \sim B(n, p)$  so that:

$$P\{Y(t) = k \mid X(t) = n\} = \binom{n}{k} p^k q^{n-k}, \quad 0 \leq k \leq n,$$

And:


$$P\{X(t) = n\} = e^{-\lambda t} \frac{(\lambda t)^n}{n!}.$$

$$\begin{aligned}
P\{Y(t) = k\} &= e^{-\lambda t} \sum_{n=k}^{\infty} \frac{n!}{(n-k)!k!} p^k q^{n-k} \frac{(\lambda t)^n}{n!} = \frac{p^k e^{-\lambda t}}{k!} (\lambda t)^k \underbrace{\sum_{n=k}^{\infty} \frac{(q\lambda t)^{n-k}}{(n-k)!}}_{e^{q\lambda t}} \\
&= (\lambda pt)^k \frac{e^{-(1-q)\lambda t}}{k!} = e^{-\lambda pt} \frac{(\lambda pt)^k}{k!}, \quad k = 0, 1, 2, \dots \\
&\sim P(\lambda pt).
\end{aligned}$$

More generally:


$$\begin{aligned}
P\{Y(t) = k, Z(t) = m\} &= P\{Y(t) = k, X(t) - Y(t) = m\} \\
&= P\{Y(t) = k, X(t) = k + m\} \\
&= P\{Y(t) = k \mid X(t) = k + m\} P\{X(t) = k + m\} \\
&= \binom{k+m}{k} p^k q^m \cdot e^{-\lambda t} \frac{(\lambda t)^{k+m}}{(k+m)!} = e^{-\lambda pt} \underbrace{\frac{(\lambda pt)^k}{k!}}_{P(Y(t)=k)} e^{-\lambda qt} \underbrace{\frac{(\lambda qt)^m}{m!}}_{P(Z(t)=m)} \\
&= P\{Y(t) = k\} P\{Z(t) = m\},
\end{aligned}$$

- Notice that  $Y(t)$  and  $Z(t)$  are generated as a result of **random Bernoulli selections** from the **original Poisson process**  $X(t)$ , where each arrival gets tossed over to either  $Y(t)$  with probability  $p$  or to  $Z(t)$  with probability  $q$ . Each such **sub-arrival** stream is also a **Poisson process**. Thus, a random selection of Poisson points preserves the Poisson nature of the resulting processes.
- However, a deterministic selection from a Poisson process destroys the Poisson property for the resulting processes.



# Inter-arrival Distribution for Poisson Processes

Let  $\tau_1$  denote the time interval (delay) to the first arrival from *any* fixed point  $t_0$ . To determine the probability distribution of the random variable  $\tau_1$ , we argue as follows: Observe that the **event** " $\tau_1 > t$ " is the same as " $n(t_0, t_0+t) = 0$ ", or the **complement event** " $\tau_1 \leq t$ " is the same as the event " $n(t_0, t_0+t) > 0$ ".



# Inter-arrival Distribution for Poisson Processes

Hence the **distribution function** of  $\tau_1$  is given by:

$$\begin{aligned} F_{\tau_1}(t) &\triangleq P\{\tau_1 \leq t\} = P\{X(t) > 0\} = P\{n(t_0, t_0 + t) > 0\} \\ &= 1 - P\{n(t_0, t_0 + t) = 0\} = 1 - e^{-\lambda t} \end{aligned}$$

Its derivative gives **the probability density function** for  $\tau_1$  to be:

$$f_{\tau_1}(t) = \frac{dF_{\tau_1}(t)}{dt} = \lambda e^{-\lambda t}, \quad t \geq 0$$

i.e.  $\tau_1$  is an exponential random variable with parameter  $\lambda$  so that:  $E(\tau_1) = 1/\lambda$ .

# Inter-arrival Distribution for Poisson Processes

Similarly, let  $t_n$  represent the  $n^{\text{th}}$  random arrival point for a Poisson process. Then:

$$\begin{aligned}\Delta F_{t_n}(t) &= P\{t_n \leq t\} = P\{X(t) \geq n\} \\ &= 1 - P\{X(t) < n\} = 1 - \sum_{k=0}^{n-1} \frac{(\lambda t)^k}{k!} e^{-\lambda t}\end{aligned}$$

and hence:

$$\begin{aligned}f_{t_n}(x) &= \frac{dF_{t_n}(x)}{dx} = -\sum_{k=1}^{n-1} \frac{\lambda(\lambda x)^{k-1}}{(k-1)!} e^{-\lambda x} + \sum_{k=0}^{n-1} \frac{\lambda(\lambda x)^k}{k!} e^{-\lambda x} \\ &= \frac{\lambda^n x^{n-1}}{(n-1)!} e^{-\lambda x}, \quad x \geq 0\end{aligned}$$

# Inter-arrival Distribution for Poisson Processes

which represents a Gamma density function. i.e., the **waiting time** to the  $n^{\text{th}}$  **Poisson arrival** has a **Gamma distribution**.

Moreover:

$$t_n = \sum_{i=1}^n \tau_i$$

where  $\tau_i$  is the random inter-arrival duration between the  $(i - 1)^{\text{th}}$  and  $i^{\text{th}}$  events. Notice that  $\tau_i$  s are **independent, identically distributed random variables**. Hence using their characteristic functions, it follows that all inter-arrival durations of a Poisson process are independent exponential random variables with common parameter  $\lambda$ .

i.e.,

$$f_{\tau_i}(t) = \lambda e^{-\lambda t}, \quad t \geq 0.$$



# Inter-arrival Distribution for Poisson Processes

Alternatively, we have  $\tau_1$  is an exponential random variable. By repeating that argument after shifting  $t_0$  to the new point  $t_1$ , we conclude that  $\tau_2$  is an exponential random variable. Thus, the sequence  $\tau_1, \tau_2, \dots, \tau_n, \dots$  are **independent exponential random variables** with common p.d.f.

Thus, if we systematically tag every  $m^{\text{th}}$  outcome of a Poisson process  $X(t)$  with parameter  $\lambda t$  to generate a new process  $e(t)$ , then the inter-arrival time between any two events of  $e(t)$  is a **Gamma random variable**.

# Inter-arrival Distribution for Poisson Processes

Notice that:


$$E[e(t)] = m / \lambda, \text{ and if } \lambda = m\mu, \text{ then } E[e(t)] = 1 / \mu.$$

The inter-arrival time of  $e(t)$  in that case represents an **Erlang- $m$  random variable**, and  $e(t)$  is an **Erlang- $m$  process**.

In summary, if Poisson arrivals are randomly redirected to form new queues, then each such queue generates a new Poisson process.

# Poisson Departures between Exponential Inter-arrivals

Let  $X(t) \sim P(\lambda t)$  and  $Y(t) \sim P(\mu t)$  represent two independent Poisson processes called *arrival* and *departure* processes.



Let  $Z$  represent the random interval between *any* two successive arrivals of  $X(t)$ .  $Z$  has an exponential distribution with parameter  $\lambda$ . Let  $N$  represent the number of “departures” of  $Y(t)$  between *any* two successive arrivals of  $X(t)$ . Then from the Poisson nature of the departures we have:

$$P\{N = k \mid Z = t\} = e^{-\mu t} \frac{(\mu t)^k}{k!}.$$

# Poisson Departures between Exponential Inter-arrivals

$$\begin{aligned}P\{N = k\} &= \int_0^{\infty} P\{N = k \mid Z = t\} f_Z(t) dt \\&= \int_0^{\infty} e^{-\mu t} \frac{(\mu t)^k}{k!} \lambda e^{-\lambda t} dt \\&= \frac{\lambda}{k!} \int_0^{\infty} (\mu t)^k e^{-(\lambda+\mu)t} dt \\&= \frac{\lambda}{\lambda+\mu} \left( \frac{\mu}{\lambda+\mu} \right)^k \underbrace{\frac{1}{k!} \int_0^{\infty} x^k e^{-x} dx}_{k!} \\&= \left( \frac{\lambda}{\lambda+\mu} \right) \left( \frac{\mu}{\lambda+\mu} \right)^k, \quad k = 0, 1, 2, \dots\end{aligned}$$

# Poisson Departures between Exponential Inter-arrivals

- The random variable  $N$  has a **geometric distribution**. Thus if customers come in and get out according to two independent Poisson processes at a counter, then the number of arrivals between any two departures has a geometric distribution.
- Similarly, the number of departures between *any* two arrivals also represents another geometric distribution.

## Example

Suppose there are 2 independent Poisson processes with  $\lambda_1 = 1, \lambda_2 = 2$ .

Find the probability that 2<sup>nd</sup> arrival of first process occurs before 3<sup>rd</sup> arrival of the second process.

### Solution:

Consider the superposition of these two Poisson processes. It is still a Poisson process with  $\lambda = 1 + 2 = 3$ .

Also, each event of the resulting process is from first process with probability  $\frac{\lambda_1}{\lambda_1 + \lambda_2} = \frac{1}{3}$  and otherwise with probability  $\frac{2}{3}$ . So, for the 2<sup>nd</sup> arrival of first process to occur before 3<sup>rd</sup> arrival of the second process, we need the first 4 occurrences to cover at least 2 occurrences of the first process:

$$\sum_{k=2}^4 \binom{4}{k} \left(\frac{1}{3}\right)^k \left(\frac{2}{3}\right)^{4-k}$$

## Example: Coupon Collecting

Suppose a cereal manufacturer randomly inserts a sample of one type of coupon into each cereal box. Suppose there are  $n$  such distinct types of coupons. One interesting question is how many boxes of cereal should one buy on average to collect at least one coupon of each kind?

## Example: Coupon Collecting

We shall reformulate the above problem in terms of Poisson processes. Let  $X_1(t), X_2(t), \dots, X_n(t)$  represent  $n$  *independent* identically distributed Poisson processes with common parameter  $\lambda t$ . Let  $t_{i1}, t_{i2}, \dots$  represent the first, second, ... random arrival instants of the process  $X_i(t)$ ,  $i = 1, 2, \dots, n$ . They will correspond to the first, second, ... appearance of the  $i^{\text{th}}$  type of coupon in the above problem. Let:

$$X(t) \triangleq \sum_{i=1}^n X_i(t),$$

so that the sum  $X(t)$  is also a Poisson process with parameter  $\mu t$ , where


$$\mu = n\lambda.$$



# Example: Coupon Collecting

$1/\lambda$  represents: The average inter-arrival duration between any two arrivals of  $X_i(t), i = 1, 2, \dots, n$ , whereas:

$1/\mu$  represents the average inter-arrival time for the combined sum process  $X(t)$ .




# Bulk Arrivals and Compound Poisson Processes


In an ordinary Poisson process  $X(t)$ , only one event occurs at any arrival instant. Instead suppose a random number of events  $C_i$  occur simultaneously as a cluster at every arrival instant of a Poisson process. If  $X(t)$  represents the total number of all occurrences in the interval  $(0, t)$ , then  $X(t)$  represents a **compound Poisson process**, or a **bulk arrival process**.

# Bulk Arrivals and Compound Poisson Processes

Inventory orders, arrivals at an airport queue, tickets purchased for a show, etc. follow this process (when things happen, they happen in a bulk, or a bunch of items are involved.)



(a) Poisson Process



(b) Compound Poisson Process

Let:

$$p_k = P\{C_i = k\}, \quad k = 0, 1, 2, \dots$$

represent the common probability mass function for the occurrence in any cluster  $C_i$ . Then the compound process  $X(t)$  satisfies:

$$X(t) = \sum_{i=1}^{N(t)} C_i,$$

where  $N(t)$  represents an ordinary Poisson process with parameter  $\lambda$ .  
Let:

$$P(z) = E\{z^{C_i}\} = \sum_{k=0}^{\infty} p_k z^k$$

represent the moment generating function associated with the cluster Statistics. Then the moment generating function of the compound Poisson process  $X(t)$  is given by:

$$\begin{aligned}
 \phi_x(z) &= \sum_{n=0}^{\infty} z^n P\{X(t) = n\} = E\{z^{X(t)}\} \\
 &= E\{E[z^{X(t)} \mid N(t) = k]\} = E\{E\{z^{\sum_{i=1}^k C_i} \mid N(t) = k\}\} \\
 &= \sum_{k=0}^{\infty} (E\{z^{C_i}\})^k P\{N(t) = k\} \\
 &= \sum_{k=0}^{\infty} P^k(z) e^{-\lambda t} \frac{(\lambda t)^k}{k!} = e^{-\lambda t(1-P(z))}
 \end{aligned}$$

If we let:

$$P^k(z) \triangleq \left( \sum_{n=0}^{\infty} p_n z^n \right)^k = \sum_{n=0}^{\infty} p_n^{(k)} z^n$$

where  $\{p_n^{(k)}\}$  represents the  $k$  fold convolution of the sequence  $\{p_n\}$  with itself, we obtain:

$$P\{X(t) = n\} = \sum_{k=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^k}{k!} p_n^{(k)}$$

The above, represents the probability that there are  $n$  arrivals in the interval  $(0, t)$  for a compound Poisson process  $X(t)$ .

We can rewrite  $\phi_X(z)$  also as:

$$\phi_X(z) = e^{-\lambda_1 t(1-z)} e^{-\lambda_2 t(1-z^2)} \dots e^{-\lambda_k t(1-z^k)} \dots$$

where  $\lambda_k = p_k \lambda$ , which shows that the compound Poisson process can be expressed as the sum of integer-scaled independent Poisson processes  $m_1(t), m_2(t), \dots$ . Thus:

$$X(t) = \sum_{k=1}^{\infty} k m_k(t).$$


More generally, every linear combination of independent Poisson processes represents a compound Poisson process.

# Outline of Week 04 Lectures


- Poisson Process
- Point Process



# Many discrete *events* in continuous time




## Online actions



## Disease dynamics



## Financial trading



## Mobility dynamics

Variety of processes behind these events



**Events are (noisy) observations of a variety of complex dynamic processes...**



**...in a wide range of temporal scales.**

# Example I: Information propagation

$S \rightarrow D$   
means  
D follows S



**They can have an impact  
in the off-line world**

**theguardian**

Click and elect: how fake news helped Donald Trump win a real election



### Barack Obama

From Wikipedia, the free encyclopedia

*"Barack" and "Obama" redirect here. For his father, see Barack Obama Sr. For other uses of "Barack", see Barack (disambiguation) (disambiguation).*

**Barack Hussein Obama II** (), current President of the United States. He was president of the Harvard Law School, a civil rights attorney and taught representing the 13th District States House of Representatives.

### Barack Obama: Revision history

- 03:41, 28 November 2016 Ranze (talk | contribs) .. (301,105 bytes) (+18) .. (E)
- 03:32, 28 November 2016 Xin Deui (talk | contribs) .. (301,087 bytes) (-68) .. (E)
- 00:57, 28 November 2016 SporkBot (talk | contribs) m .. (301,155 bytes) (-37) .. (E)
- 07:03, 27 November 2016 Saiph121 (talk | contribs) .. (301,192 bytes) (+25) .. (E)

03:21, 20 September 2016



is a **Kenyan** politician



possible vandalism by **MLM2016**

is an American politician

- Addition
- Refutation



Moving to Australia Working in Australia Study abroad in Australia +4

### What are the pros and cons of living in Australia?

Answer Request Follow 109 Comment Share 9 Downvote

I have studied, worked and lived in Australia as an Internat employee, business owner and a citizen.

Upvote | 150

I have experienced this country in all the ways possible, you However, I firmly believe that there are definitely more pros Australia but still I have mentioned below a few challenges and benefits.





**M Sharma**, Lived in Australia as Migrant, Student, Worker, Business Owner & Family Man

Updated Aug 3



Hope it helps! :)

Possible Challenges

- Language problem for those who don't speak English
- Not having your family and friends around could be a challenge. However, the Australian society is more and more connected and thanks to Social Media you can stay in touch a bit easier with your family and friends.




# Aren't these event traces just time series?



**Discrete and continuous times series**

**Discrete events in continuous time**

**What about aggregating events in *epochs*?**



- How long is each epoch?
- How to aggregate events per epoch?
- What if no event in one epoch?
- What about time-related queries?


# Temporal Point Processes (TPPs): Introduction

- 1. Intensity function**
2. Basic building blocks
3. Superposition
4. Marks and SDEs with jumps

# Temporal point processes

## Temporal point process:



A random process whose realization consists of discrete events localized in time  $\mathcal{H} = \{t_i\}$



**Formally:**  $N(t) = \int_0^t dN(s)$   $\Rightarrow$   $dN(t) \in \{0, 1\}$   $\Downarrow$   $dN(t) = \sum_{t_i \in \mathcal{H}} \delta(t - t_i) dt$

Dirac delta function  $\Downarrow$


# Model time as a random variable



Likelihood of a timeline:  $f^*(t_1) f^*(t_2) f^*(t_3) f^*(t) S^*(T)$



# Problems of density parametrization (I)




$$\begin{array}{cccccc}
 f^*(t_1) & f^*(t_2) & f^*(t_3) & f^*(t) & S^*(T) & \\
 \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \\
 \frac{\exp\langle w, \psi^*(t_1) \rangle}{Z} & \frac{\exp\langle w, \psi^*(t_2) \rangle}{Z} & \frac{\exp\langle w, \psi^*(t_3) \rangle}{Z} & \frac{\exp\langle w, \psi^*(t) \rangle}{Z} & 1 - \int_t^T \frac{\exp\langle w, \psi^*(\tau) \rangle}{Z} d\tau & 
 \end{array}$$

It is **difficult for model design and interpretability**:

1. Densities need to integrate to 1 (i.e., partition function)
2. Difficult to combine timelines

# Intensity function




**Intensity:**

Probability between  $[t, t+dt)$  but not before  $t$

$$\lambda^*(t) dt = \frac{f^*(t) dt}{S^*(t)} \geq 0 \quad \Rightarrow \quad \lambda^*(t) dt = \mathbb{E}[dN(t) | \mathcal{H}(t)]$$

**Observation:**  $\lambda^*(t)$  It is a rate = # of events / unit of time

# Advantages of intensity parametrization (I)



$$\lambda^*(t_1) \lambda^*(t_2) \lambda^*(t_3) \lambda^*(t) \exp\left(-\int_0^T \lambda^*(\tau) d\tau\right)$$


$$\langle w, \phi^*(t_1) \rangle \quad \langle w, \phi^*(t_2) \rangle \quad \langle w, \phi^*(t_3) \rangle \quad \langle w, \phi^*(t) \rangle \quad \exp\left(-\int_0^T \langle w, \phi^*(\tau) \rangle d\tau\right)$$

Arrows point from the inner product terms to the corresponding  $\lambda^*$  terms in the first equation, and from the exponential term to the corresponding exponential term in the second equation.

**Suitable for model design and interpretable:**

1. Intensities only need to be nonnegative
2. Easy to combine timelines

# Relation between $f^*$ , $F^*$ , $S^*$ , $\lambda^*$



# Representation:

## Temporal Point Processes

1. Intensity function
- 2. Basic building blocks**
3. Superposition
4. Marks and SDEs with jumps


## Recall: Some Sampling Techniques

- Sampling is essential in statistics because it makes inference more efficient, feasible, accurate, and resource-effective while allowing for generalizability and detailed analysis.
- We treat sampling methods in more detail at the end of the course.
- **Inversion sampling:** Also known as inverse transform sampling, is a method for generating random samples from any probability distribution given its cumulative distribution function (CDF), in two steps:
  - Uniform Random Sample: Generate a random number ( $u$ ) from a uniform distribution between 0 and 1.
  - Inverse CDF: Use the inverse of the cumulative distribution function (CDF) of the target distribution to transform the uniform random sample. This involves finding the value ( $x$ ) such that  $(F(x) = u)$ , where ( $F$ ) is the CDF of the target distribution.

## Recall: Some Sampling Techniques

- **Rejection sampling:** also known as the acceptance-rejection method, is a technique used in computational statistics to generate observations from a target distribution by using a proposal distribution:
  - **Proposal Distribution:** Choose a proposal distribution ( $g(x)$ ) from which it is easy to sample. This distribution should cover the support of the target distribution ( $f(x)$ ).
  - **Sampling:** Generate a samples ( $x$ ) from the proposal distribution ( $g(x)$ ).
  - **Acceptance Criterion:** Accept the sample ( $x$ ) if the defined acceptance criterion is met. Repeat the process until a sample is accepted.

# Poisson process



## Intensity of a Poisson process


$$\lambda^*(t) = \mu$$

### Observations:

1. Intensity independent of history
2. Uniformly random occurrence
3. Time interval follows exponential distribution



# Fitting & sampling from a Poisson



**Fitting by maximum likelihood:**


$$\mu^* = \operatorname{argmax}_{\mu} 3 \log \mu - \mu T = \frac{3}{T}$$

**Sampling using inversion sampling:**

$$t \sim \underbrace{\mu \exp(-\mu(t - t_3))}_{f_t^*(t)} \quad \rightarrow \quad t = \underbrace{-\frac{1}{\mu} \log(1 - u)}_{F_t^{-1}(u)} + t_3$$

*Uniform(0, 1)*  
↓


# Inhomogeneous Poisson process




## Intensity of an inhomogeneous Poisson process

$$\lambda^*(t) = g(t) \geq 0 \quad (\text{Independent of history})$$

Example:



# Fitting & sampling from inhomogeneous Poisson




**Fitting by maximum likelihood:**  $\underset{g(t)}{\text{maximize}} \sum_{i=1}^n \log g(t_i) - \int_0^T g(\tau) d\tau.$

**Sampling using thinning (reject. sampling) + inverse sampling:**

1. Sample  $t$  from Poisson process with intensity  $\mu$  using inverse sampling
  2. Generate  $u_2 \sim \text{Uniform}(0, 1)$
  3. Keep the sample if  $u_2 \leq g(t) / \mu$
- } Keep sample with prob.  $g(t) / \mu$

# Terminating (or survival) process



## Intensity of a terminating (or survival) process


$$\lambda^*(t) = g^*(t)(1 - N(t)) \geq 0$$

### Observations:

1. Limited number of occurrences

*Try sampling  
and fitting!*

# Self-exciting (or Hawkes) process




Intensity of self-exciting  
(or Hawkes) process:

$$\begin{aligned}\lambda^*(t) &= \mu + \alpha \sum_{t_i \in \mathcal{H}(t)} \kappa_\omega(t - t_i) \\ &= \mu + \alpha \kappa_\omega(t) \star dN(t)\end{aligned}$$

Observations:

1. Clustered (or bursty) occurrence of events
2. Intensity is stochastic and history dependent

# Fitting a Hawkes process from a recorded timeline



## Fitting by maximum likelihood:

$$\text{maximize}_{\mu, \alpha} \left. \sum_{i=1}^n \log \lambda^*(t_i) - \int_0^T \lambda^*(\tau) d\tau \right\} \begin{array}{l} \text{The max. likelihood} \\ \text{is jointly convex} \\ \text{in } \mu \text{ and } \alpha \end{array}$$

## Sampling using thinning (reject. sampling) + inverse sampling:


Key idea: the maximum of the intensity  $\lambda_0$  changes over time

# Summary

## Building blocks to represent different dynamic processes:


Poisson processes:

$$\lambda^*(t) = \lambda$$




Inhomogeneous Poisson processes:

$$\lambda^*(t) = g(t)$$




Terminating point processes:

$$\lambda^*(t) = g^*(t)(1 - N(t))$$



Self-exciting point processes:

$$\lambda^*(t) = \mu + \alpha \sum_{t_i \in \mathcal{H}(t)} \kappa_\omega(t - t_i)$$




# Representation:

## Temporal Point Processes

1. Intensity function
2. Basic building blocks
- 3. Superposition**
4. Marks and SDEs with jumps




# Mutually exciting process



## Clustered occurrence affected by neighbors

$$\lambda^*(t) = \mu + \alpha \sum_{t_i \in \mathcal{H}_b(t)} \kappa_\omega(t - t_i) + \beta \sum_{t_i \in \mathcal{H}_c(t)} \kappa_\omega(t - t_i)$$

# Mutually exciting terminating process



## Clustered occurrence affected by neighbors

$$\lambda^*(t) = (1 - N(t)) \left( g(t) + \beta \sum_{t_i \in \mathcal{H}_c(t)} \kappa_\omega(t - t_i) \right)$$

# Representation:


## Temporal Point Processes

1. Intensity function
2. Basic building blocks
3. Superposition
- 4. Marks and SDEs with jumps**


# Marked temporal point processes

## Marked temporal point process:

A random process whose realization consists of **discrete marked events localized in time**



# Independent identically distributed marks




**Distribution for the marks:**

$$x^*(t_i) \sim p(x)$$

**Observations:**

1. Marks independent of the temporal dynamics
2. Independent identically distributed (I.I.D.)

# Dependent marks: SDEs with jumps




Marks given by stochastic differential equation with jumps:

$$x(t + dt) - x(t) = dx(t) = \underbrace{f(x(t), t)dt}_{\text{Drift}} + \underbrace{h(x(t), t)dN(t)}_{\text{Event influence}}$$

Observations:

1. Marks dependent of the temporal dynamics
2. Defined for all values of  $t$

# Dependent marks: distribution + SDE with jumps




Distribution for the marks:

$$x^*(t_i) \sim p(x^* | x(t)) \Rightarrow dx(t) = \underbrace{f(x(t), t)dt}_{\text{Drift}} + \underbrace{h(x(t), t)dN(t)}_{\text{Event influence}}$$

Observations:

1. Marks dependent on the temporal dynamics
2. Distribution represents additional source of uncertainty

# Mutually exciting + marks




## Marks affected by neighbors

$$dx(t) = \underbrace{f(x(t), t)dt}_{\text{Drift}} + \underbrace{g(x(t), t)dM(t)}_{\text{Neighbor influence}}$$




# Marked TPPs as stochastic dynamical systems

## Example: Susceptible-Infected-Susceptible (SIS)




SDE with jumps

$$dX_i(t) = \underbrace{dY_i(t)}_{\text{It gets infected}} - \underbrace{dW_i(t)}_{\text{It recovers}}$$



Node is susceptible

$$\lambda_{Y_i}(t)dt = (1 - X_i(t))\beta \underbrace{\sum_{j \in \mathcal{N}(i)} X_j(t)}_{\text{If friends are infected, higher infection rate}} dt$$



SDE with jumps


$$d\lambda_{W_i}(t) = \underbrace{\delta dY_i(t)}_{\text{Self-recovery rate when node gets infected}} - \underbrace{\lambda_{W_i}(t)dW_i(t)}_{\text{If node recovers, rate to zero}} + \underbrace{\rho dN_i(t)}_{\text{Rate increases if node gets treated}}$$

# Models & Inference


- 1. Modeling event sequences**
2. Clustering event sequences
3. Capturing complex dynamics
4. Causal reasoning on event sequences

# Event sequences as cascades

$S \rightarrow D$   
S means  
D follows S





## Information Diffusion



[Leskovec et al., 2009]



## Disease Diffusion



[Rizoiu et al., 2018]

# An example: idea adoption

**S** → **D**  
means  
D follows S




**They can have an impact  
in the off-line world**

**theguardian**


Click and elect: how fake news helped Donald Trump win a real election

# Infection cascade representation


We represent an infection cascade using **terminating temporal point processes**:



**Infection event:**



# Infection intensity




$$\lambda_u^*(t) = \underbrace{(1 - N_u(t))}_{\text{Users get infected only once}} \sum_{v \in [m]} \underbrace{b_{vu} \sum_{e_i \in \mathcal{H}_v(t)} \kappa(t - t_i)}_{\text{Influence from user } v \text{ on user } u \text{ Previous infections of user } v}$$

# Model inference from multiple cascades

Conditional intensities

$$\lambda_u^*(t)$$



Diffusion log-likelihood

$$\mathcal{L} = \sum_{u=1}^n \log \lambda_u^*(t_u) - \int_0^T \lambda_u^*(\tau) d\tau$$

Maximum likelihood approach to find model parameters!




Sum up log-likelihoods of multiple cascades!

**Theorem.** For any choice of parametric memory, the **maximum likelihood** problem is **convex**.

In some cases, influence change over time:



Propagation over networks  
with variable influence






# Recurrent events: beyond cascades

**Up to this point**, each users is only infected once, and event sequences can be seen as cascades.

**In general, users perform recurrent events over time.** E.g., people repeatedly express their opinion online:



How social media is revolutionizing debates

*The New York Times*

*Social Media Are Giving a Voice to Taste Buds*




**Twitter Unveils A New Set Of Brand-Centric Analytics**

*The New York Times*


*Campaigns Use Social Media to Lure Younger Voters*

# Recurrent events representation


We represent messages using **nonterminating temporal point processes**:



**Recurrent event:**




# Recurrent events intensity



**Cascade sources!**

$$\underbrace{\lambda_u^*(t)}_{\text{User's intensity}} = \underbrace{\mu_u}_{\text{Events on her own initiative}} + \sum_{v \in [m]} \underbrace{b_{vu} \sum_{e_i \in \mathcal{H}_v(t)} \kappa(t - t_i)}_{\text{Influence from user } v \text{ on user } u}$$




**Hawkes process**

# Models & Inference

1. Modeling event sequences
- 2. Clustering event sequences**
3. Capturing complex dynamics
4. Causal reasoning on event sequences


# Event sequences

So far, we have assumed the cascade (topic, meme, etc.) that each event belongs to was known.




Often, the cluster (topic, meme, etc.) that each event in a sequence belongs to is not known:


-  **BBC News (World)** @BBCWorld · 4m  
Turkey election: Erdogan win ushers in new presidential era
-  **BBC News (World)** @BBCWorld · 46m  
Dublin church: Seven injured as car hits pedestrians
-  **BBC News (World)** @BBCWorld · 2h  
Nigerian music star D'banj's son 'drowns at home'
-  **BBC News (World)** @BBCWorld · 2h  
Turkey election: Country's heart split over Erdogan victory



Assume the event **cluster** to be **hidden** and aim to automatically **learn the cluster assignments** from the data:




**Bayesian methods** to cluster event sequences in the context of:



**Learning**

### Online News



### Health care

Method	DMHP
ICU Patient	<b>0.3778</b>
IPTV User	<b>0.2004</b>


[Du et al., 2015; Mavroforakis et al., 2017; Xu & Zha, 2017]

# Hierarchical Dirichlet Hawkes process




## 1st year computer science student

- Introduction to programming
- Discrete math
- Project presentation






# Events representation

We represent the events using **marked temporal point processes**:





# Cluster intensity



New cascade rate


Cluster popularity

$$\underbrace{\lambda_{u,l}^*(t)}_{\substack{\text{Intensity} \\ \text{or rate} \\ \text{(events / hour)}}} = \underbrace{\mu_u \pi_l}_{\text{Own initiative}} + \underbrace{\sum_{j:t_j \in \mathcal{H}_{u,l}(t)} k_{\theta_l}(t - t_j)}_{\text{Follow-up}}$$

**Hawkes process**

# User events intensity

Users adopt more than one cluster:




A user's learning events as a multidimensional Hawkes:

$$\begin{array}{l} \text{Time} \searrow \\ \text{cluster} \swarrow \\ (t_n, p_n) \sim \text{Hawkes} \end{array} \begin{pmatrix} \lambda_{u,1}^*(t) \\ \vdots \\ \lambda_{u,\infty}^*(t) \end{pmatrix}$$

$$\text{Content} \rightarrow q_n = \omega \quad \omega_j \sim \text{Multinomial}(\theta_p)$$

# People share same clusters

*Different users adopt same clusters*



Cluster distribution from a **Dirichlet process**:


- Infinite # of clusters.
- Shared parameters across users.

**Details in the reference below!**

# Content




# Intensities




**Version control tasks tend to be specific,  
quickly solved after performing few questions**

# Content



# Intensities



**Machine learning tasks tend to be more complex and require asking more questions**

# Models & Inference


1. Modeling event sequences
2. Clustering event sequences
- 3. Capturing complex dynamics**
4. Causal reasoning on event sequences

# **Case Studies & References**

For those who want to do research in social media


Up to now, we have focused on simple temporal dynamics (and intensity functions):

$$\lambda^*(t) = \mu$$



$$\lambda^*(t) = \sum_j \alpha_j k(t - t_j)$$

$$\lambda^*(t) = \mu + \alpha \sum_{t_i \in \mathcal{H}(t)} \kappa_\omega(t - t_i)$$




Recent works make use of **RNNs** to capture more complex dynamics

[Du et al., 2016; Dai et al., 2016; Mei & Eisner, 2017; Jing & Smola, 2017; Trivedi et al., 2017; Xiao et al., 2017a; 2018]



# Neural Hawkes process

- 1) History effect does not need to be additive
- 2) Allows for complex memory effects (such as delays)




# Neural Hawkes process

$$\lambda_u(t) = f_u(\mathbf{w}_u^\top \mathbf{h}(t))$$


Excitation & inhibition

$$\mathbf{h}(t) = \overbrace{\text{RNN}(\mathcal{H}(t))}^{\text{Memory}}$$




# Applications (I): Predictive Models


## Know-Evolve, Trivedi et al. (2017)




## Coevolutionary Embedding, Dai et al. (2017)



## Neural Survival Recommender, Jing & Smola (2017)




# Key idea: Intensity- and likelihood-free models



## Wasserstein-Distance for Temporal Point Processes

### GAN architecture






# Models & Inference



1. Modeling event sequences
2. Clustering event sequences
3. Capturing complex dynamics
- 4. Causal reasoning on event sequences**

# Temporal point processes beyond prediction

So far, we have focused on models that improve predictions:



Recent works have focused on performing **causal inference** using event sequences:




# Uncovering Causality from Hawkes Processes

## Multivariate Hawkes process:

$$N(t) = \sum_{u \in \mathcal{U}} N_u(t)$$

$$\lambda_u(t) = \mu_u + \sum_{v \in \mathcal{U}} \underbrace{\int_0^t k_{u,v}(t-t') dN_v(t')}_{\text{Effect of } v\text{'s past events on } u}$$



## Granger causality:

"X causes Y in the sense of Granger causality if forecasting future values of Y is more successful while taking X past values into account"


[Granger, 1969]

# Uncovering Causality from Hawkes Processes

## Multivariate Hawkes process:

$$N(t) = \sum_{u \in \mathcal{U}} N_u(t)$$

$$\lambda_u(t) = \mu_u + \sum_{v \in \mathcal{U}} \underbrace{\int_0^t k_{u,v}(t-t') dN_v(t')}_{\text{Effect of } v\text{'s past events on } u}$$



## Granger causality on multivariate Hawkes processes:

“  $N_v(t)$  does not Granger-cause  $N_u(t)$  w.r.t.  $N(t)$  if and only if  $k_{u,v}(\tau) = 0$  for  $\tau \in \mathbb{R}^+$  ”

[Eichler et al., 2016]

[Achab et al., ICML 2017]




# Uncovering Causality from Hawkes Processes

Goal is to estimate  $G = [g_{uv}]$ , where:

$$g_{uv} = \int_0^{+\infty} k_{u,v}(\tau) d\tau \geq 0 \text{ for all } u, v \in \mathcal{U}$$

Average total # of events of node  $u$  whose *direct* ancestor is an event by node  $v$




Then,  $G = [g_{uv}]$  quantifies the *direct causal relationship* between nodes.

# Uncovering Causality from Hawkes Processes

Goal is to estimate  $G = [g_{uv}]$ , where:

$$g_{uv} = \int_0^{+\infty} k_{u,v}(\tau) d\tau \geq 0 \text{ for all } u, v \in \mathcal{U}$$

Average total # of events of node  $u$  whose *direct* ancestor is an event by node  $v$



Then,  $G = [g_{uv}]$  quantifies the *direct causal relationship* between nodes.


**Key idea:** Estimate  $G$  using the cumulants  $dN(t)$  of the Hawkes process.

# Uncovering Causality from Hawkes Processes

Goal is to estimate  $G = [g_{uv}]$ , where:

$$g_{uv} = \int_0^{+\infty} k_{u,v}(\tau) d\tau \geq 0 \text{ for all } u, v \in \mathcal{U}$$

Average total # of events of node  $u$  whose *direct* ancestor is an event by node  $v$



Then,  $G = [g_{uv}]$  quantifies the *direct causal relationship* between nodes.

**Details in the reference below!**

**Key idea:** Estimate  $G$  using the cumulants the  $dN(t)$  of the Hawkes process.

**Next Week:**

**Gaussian Process**

**Have a good day!**